

Equivalent topological invariants of topological insulators

Zhong Wang^{1,2}, Xiao-Liang Qi^{2,3} and Shou-Cheng Zhang²

¹ Department of Modern Physics, University of Science and Technology of China, Hefei, 230026, P. R. China

² Department of Physics, Stanford University, Stanford, CA 94305, USA

³ Microsoft Research, Station Q, Elings Hall, University of California, Santa Barbara, CA 93106, USA

E-mail: sczhang@stanford.edu

Abstract. A time-reversal invariant topological insulator can be generally defined by the effective topological field theory with a quantized θ coefficient, which can only take values of 0 or π . This theory is generally valid for an arbitrarily interacting system and the quantization of the θ invariant can be directly measured experimentally. Reduced to the case of a non-interacting system, the θ invariant can be expressed as an integral over the entire three dimensional Brillouin zone. Alternatively, non-interacting insulators can be classified by topological invariants defined over discrete time-reversal invariant momenta. In this paper, we show the complete equivalence between the integral and the discrete invariants of the topological insulator.

1. Introduction

In condensed matter systems, most states of matter are classified by the symmetries they break. For example, a crystal breaks the translational symmetry, a magnet breaks the rotational symmetry and a superconductor breaks the gauge symmetry. These broken symmetry states are identified by the order parameter, and described by the effective field theory. The effective field theory is constrained by the broken symmetry, and it defines universality classes of different states of matter, and predicts universal physical properties.

The quantum Hall (QH) state is a topological state of matter which does not fall into the conventional Landau paradigm. The long distance and low energy properties of the QH state is generally described by the effective topological field theory

$$S_{\text{eff}} = \frac{C_1}{4\pi} \int d^2x dt \epsilon^{\mu\nu\tau} A_\mu \partial_\nu A_\tau, \quad (1)$$

This topological field theory is generally valid for interacting systems, and describes the electromagnetic response in the long wave length limit[1]. In the non-interacting limit, the Hall conductance $\sigma_{xy} = C_1$ can be expressed as a topological invariant over the two-dimensional (2D) Brillouin zone[2], given by

$$C_1 = \frac{e^2}{h} \frac{1}{2\pi} \int dk_x \int dk_y f_{xy}(\mathbf{k}) \quad (2)$$

where

$$f_{xy}(\mathbf{k}) = \frac{\partial a_y(\mathbf{k})}{\partial k_x} - \frac{\partial a_x(\mathbf{k})}{\partial k_y}$$

$$a_i(\mathbf{k}) = -i \sum_{\alpha \in \text{occ}} \langle \alpha \mathbf{k} | \frac{\partial}{\partial k_i} | \alpha \mathbf{k} \rangle, \quad i = x, y.$$

are the abelian Berry curvature and potential for the band states $|\alpha \mathbf{k}\rangle$. Time reversal (TR) symmetry breaking is essential for the QH effect.

More recently, new topological insulator states have been theoretically predicted and experimentally observed in HgTe quantum wells, BiSb alloys, Bi₂Te₃ and Bi₂Se₃ crystals[3; 4; 5; 6; 7; 8; 9]. These topological insulator states physically arises from the spin-orbit coupling in electronic structures, and are protected by the TR symmetry[5; 10; 11; 12; 13]. A general theory describes the three dimensional (3D) topological insulator in terms of a effective topological field theory[12], given by

$$S_{\text{eff}} = S_{\text{Maxwell}} + S_{\text{topo}} = \int d^3x dt \left[\frac{1}{16\pi} F_{\mu\nu} F^{\mu\nu} + \frac{\theta\alpha}{32\pi^2} \epsilon^{\mu\nu\sigma\tau} F_{\mu\nu} F_{\sigma\tau} \right] \quad (3)$$

where the last term is the topological term and $\alpha = e^2/hc$ is the fine structure constant. For a period system, the partition function is invariant under the shift of θ by integer multiples of 2π . Since TR transformation replaces θ by $-\theta$, only two special, quantized values of $\theta = 0$ and $\theta = \pi$ are consistent with the TR symmetry. Therefore, all TR invariant insulators fall into the two distinct classes, disconnected from each other. The

Standard Model of elementary particles also admits such a topological term, therefore, the study of the topological insulator can also shed light on the fundamental topological interactions in Nature[14].

We can always couple an arbitrarily interacting electron system to the external electromagnetic field, and integrate out all the electronic degrees of freedom to obtain the effective field theory (3). For a TR invariant insulator, only two discrete possibilities of $\theta = 0$ and $\theta = \pi$ can be realized. For a non-interacting band insulator, Qi, Hughes and Zhang (QHZ) gave an explicit formula for the θ parameter as an integral over the 3D Brillouin zone

$$\theta \equiv 2\pi P_3(\theta) = \frac{1}{16\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} \{ [f_{ij}(\mathbf{k}) - \frac{2}{3} i a_i(\mathbf{k}) \cdot a_j(\mathbf{k})] \cdot a_k(\mathbf{k}) \} \quad (4)$$

where

$$\begin{aligned} f_{ij}^{\alpha\beta} &= \partial_i a_j^{\alpha\beta} - \partial_j a_i^{\alpha\beta} + i [a_i, a_j]^{\alpha\beta}, \\ a_i^{\alpha\beta}(\mathbf{k}) &= -i \langle \alpha, \mathbf{k} | \frac{\partial}{\partial k_i} | \beta, \mathbf{k} \rangle \end{aligned}$$

are the non-abelian Berry curvature and potential for the band state $|\beta, \mathbf{k}\rangle$. This topological invariant has the physical interpretation of a magneto-electric polarization, which can be directly measured experimentally[12; 15].

We notice a beautiful symmetry between the topological field theory of the TR breaking QH state and the TR invariant topological insulator state. Both the space-time integrals (1), (3) and the Brillouin zone integrals (2), (4) are integral topological invariants in the theory of differential geometry. We see that the TR breaking QH state is described by a Chern-Simons integral over the $2+1$ dimensional space-time (1), and a first Chern integral over the 2D Brillouin zone (2), while the TR invariant topological insulator is described by the second Chern integral over the $3+1$ dimensional space-time (3) and a Chern-Simons integral (4) over the 3D Brillouin zone. These are the deepest and most natural topological invariants in mathematics, and it is gratifying to see that they also describe topological states realized in Nature.

TR invariant insulators form an universality class extending over the 4D, 3D and 2D space. The root state of this universality class is the topological insulator in 4D[16; 17]. In fact, it was the first TR invariant insulator state introduced theoretically, and historically it was referred to as the 4D QH state. It is described by a topological field theory

$$S_{\text{eff}} = \frac{C_2}{24\pi^2} \int d^4x dt \epsilon^{\mu\nu\rho\sigma\tau} A_\mu \partial_\nu A_\rho \partial_\sigma A_\tau \quad (5)$$

Under TR transformation,

$$A_0 \rightarrow A_0, \quad A_i \rightarrow -A_i \quad (6)$$

therefore, we see that the topological field theory in $2+1$ dimensions (1) breaks TR symmetry, whereas the topological field theory in $4+1$ dimensions (5) preserves the TR symmetry, and naturally describes the TR invariant topological insulators. For the

case of non-interacting fermions, the coefficient is given explicitly by the second Chern number

$$C_2 = \frac{1}{32\pi^2} \int d^4k \epsilon^{ijkl} \text{tr} [f_{ij} f_{kl}] \quad (7)$$

The TR invariant topological insulator defined by (5) and (7) in the 4D space naturally generalizes the corresponding quantities of the TR breaking topological insulator defined by (1) and (2) in the 2D space. This is the reason why it was historically referred to as the 4D QH state[16; 17]. It is the root state for all TR invariant topological insulators in 3D and 2D, which can be obtained from the root state in 4D through the process of dimensional reduction[12].

Topological invariants in 3D band insulators have been studied from a different approach[10; 13]. In a beautiful series of papers, Fu, Kane and Mele (FKM) introduced a Z_2 topological invariant[5; 10; 18] for the strong topological insulator, expressed as a discrete product over the eight time-reversal invariant momenta (TRIM), explicitly written as

$$(-1)^{\nu_0} = \prod_{i=1}^8 \delta_i \quad (8)$$

where δ_i is quantity defined at TRIM Γ_i , the detailed form of which will be given in the following sections. Since weak topological insulators may not be generally robust, we shall not be concerned with their definitions here.

Therefore, there are now two topological invariants defined for the topological insulator. They are motivated by different logical reasoning and have different mathematical forms. The integral invariant (4) given by QHZ is physically measurable in terms of the magneto-electric polarization, and leads directly to the general topological field theory (3). The discrete invariant (8) given by FKM has the distinct advantage that it can be easily evaluated, especially for crystals with inversion symmetry. Applied to concrete models of topological insulators[12; 19], these two definitions yield the same result. However, it is highly desirable to prove the general equivalence between these two definitions. In this paper, we show explicitly that one can transform the integral invariant (4) to the discrete invariant (8) exactly, proving the precise equivalence between these two definitions.

2. A pedagogical example

Because our approach involved the mathematical concept of degree of map, we shall give a brief introduction to it. Our presentation in this section is mainly designed for intuitive understanding, rather than mathematical rigor. Therefore, we shall illustrate the idea of degree of map in a simple one-dimensional example. For more mathematical details, *c. f.* Ref. [20]. Those readers who are already familiar with this subject may skip this section.

Let us consider one concrete example. Consider a map $f : M \rightarrow N$, where M and N are both one-dimensional circle S^1 (see Fig.1). The coordinates of M and N are




Figure 1. Map between two circles M and N . The arrows indicate the map. The winding number of (a), (b) and (c) is 1, 2 and 1 respectively.

denoted as ϕ and θ , respectively. The degree of map f is just the number of times that M covers N under f and it is often called “winding number”.

The standard integral form of the winding number is defined as

$$\deg(f) = \frac{1}{2\pi} \int_{\phi=0}^{\phi=2\pi} d\theta(\phi) = \frac{1}{2\pi} \int_{\phi=0}^{\phi=2\pi} \frac{d\theta}{d\phi} d\phi = n \in \mathbb{Z} \quad (9)$$

which has a simple geometrical interpretation. For simplicity, let us assume that the point $\phi = 0$ maps to $\theta = 0$. When ϕ goes from 0 to 2π , θ goes from 0 to $2\pi n$. This integral form of the winding number can be expressed in a discrete form. We arbitrarily choose a image point p on the image manifold N (see Fig.1 for illustration), and count the number of source points $\bar{p}_1, \bar{p}_2, \bar{p}_3$, etc on the source manifold M , where the source points are weighted by the $+1$ or -1 sign depending on the orientation of the map. For example, image point q has only one source point $\bar{q}$, which maps onto q in a clockwise sense. Therefore $\deg(f) = 1$. On the other hand, the image point p has three source points $\bar{p}_1, \bar{p}_2$ and $\bar{p}_3$. $\bar{p}_1$ and $\bar{p}_3$ map onto p in a clockwise sense, while $\bar{p}_2$ maps onto p in a counterclockwise sense, giving $\deg(f) = +1 - 1 + 1 = 1$. We see from this example that the integral invariant of the winding number can be reduced to counting the weighted number of source points of a given image point – this is a discrete invariant for the winding number.

The above picture can be generalized to higher dimensions. Generally, for a map $f : M \rightarrow N$, where M and N are both d dimensional orientable manifold, we can defined the degree of map as

$$\deg(f) = \int_M f^*(\omega) \quad (10)$$

where ω is a d -form volume element on N satisfying $\int_N \omega = 1$, and $f^*(\omega)$ is the pullback of ω to M under f . In our one-dimensional example discussed earlier, $\omega = d\theta/2\pi$, and $f^*(\omega) = d\theta(\phi)/2\pi = (d\theta/d\phi)d\phi/2\pi$.

In our one dimensional example, we have seen that there is a discrete form of the

invariant $\deg(f)$. This can also be generalized to higher dimension. It is given as[20]

$$\deg(f) = N[f^{-1}(p), J_{f^{-1}(p)} > 0] - N[f^{-1}(p), J_{f^{-1}(p)} < 0] \quad (11)$$

where J is the Jacobian of the map, and $N[f^{-1}(p), J_{f^{-1}(p)} > (<)0]$ denotes the number of source points mapping to p with a positive (negative) Jacobian, respectively. Without going into the general derivation of this formula, we just give the explanation in our one dimensional example. In that example, $J_{f^{-1}(p)}$ is the direction sense (clockwise or counterclockwise) of the map at the point p , which is exactly what we have discussed earlier. We also mention that the point p in Eq.(11) should be regular, which means that the Jacobian of map are nonzero at $f^{-1}(p)$. Because the set of non-regular points has zero measure, we can always perturb the map to remove non-regularity at a given point.

In summary, we showed that the degree of a map can be expressed in two equivalent forms, the integral form (10) and the discrete form (11). In the next section, we will apply this idea to the integral and the discrete invariants of (4) and (8), which are degree of map $\deg(f)$ modulo 2, simply denoted as $\deg_2(f)$.

3. Equivalence between integral and discrete topological invariants

Let us start from the band structure of time-reversal invariant topological insulators. To simplify the problem, we first assume that there are no degeneracies except those required by time reversal symmetry. Suppose that there are $2N$ filled bands labeled by $(\alpha, \beta, \dots)$, where the number of filled bands is even because of the Kramers degeneracy. The $2N$ by $2N$ matrix $B(\mathbf{k})$ is defined by

$$|-\mathbf{k}, \alpha\rangle = \sum_{\beta} B_{\alpha\beta}^*(\mathbf{k}) |\Theta, \mathbf{k}, \beta\rangle \quad (12)$$

where $|\Theta, \mathbf{k}, \beta\rangle = \hat{\Theta}|\mathbf{k}, \beta\rangle$ and $\hat{\Theta}$ is the TR operator. The property $\hat{\Theta}^2 = -1$ is crucial for TR invariant Fermi systems. By direct calculation we get the inner product

$$\begin{aligned} \langle -\mathbf{k}, \alpha | \Theta, \mathbf{k}, \beta \rangle &= \sum_{\gamma} B_{\alpha\gamma}(\mathbf{k}) \langle \Theta, \mathbf{k}, \gamma | \Theta, \mathbf{k}, \beta \rangle \\ &= \sum_{\gamma} B_{\alpha\gamma}(\mathbf{k}) \delta_{\gamma\beta} \\ &= B_{\alpha\beta}(\mathbf{k}) \end{aligned} \quad (13)$$

and an important property of $B(\mathbf{k})$

$$\begin{aligned} B_{\alpha\beta}(-\mathbf{k}) &= \langle \mathbf{k}, \alpha | \Theta, -\mathbf{k}, \beta \rangle \\ &= -\langle -\mathbf{k}, \beta | \Theta, \mathbf{k}, \alpha \rangle \\ &= -B_{\beta\alpha}(\mathbf{k}) \end{aligned} \quad (14)$$

where the facts that $\hat{\Theta}^2 = -1$ and $\langle \alpha | \Theta, \beta \rangle = -\langle \beta | \Theta, \alpha \rangle$ have been used. Therefore, $B_{\alpha\beta}(-\mathbf{k})$ is anti-symmetric at the eight TR invariant momenta(TRIM), which enables

the definition of Pfaffian at these points. The definition of the discrete invariant given by FKM[10; 18] is expressed as

$$(-1)^{\nu_0} = \prod_{i=1}^8 \delta_i \quad ; \quad \delta_i = \frac{\sqrt{\det[B(\Gamma_i)]}}{\text{Pf}[B(\Gamma_i)]} = \pm 1$$

with the Pfaffian of B written as $\text{Pf}[B]$. Although it seems that all the quantities appearing in this definition are local, global information on T^3 is encoded because a global basis of wavefunction is required in this definition. The existence of global basis is not obvious and we shall present a discussion on this in the appendix. The unitary matrix $B(\mathbf{k})$ defines a map

$$f : T^3 \rightarrow U(2N) \quad (15)$$

This map can be simplified since we have the assumption of absence of accidental degeneracies. We divide the $2N$ filled bands into N TR pairs. Because energy eigenstates with different eigenenergy are orthogonal, we have $B_{\alpha\beta}(\mathbf{k}) = 0$ if $|\mathbf{k}, \alpha\rangle$ and $|\mathbf{k}, \beta\rangle$ belong to different pairs. Therefore, all the inter-pairs elements of matrix $B(\mathbf{k})$ are zero. Written explicitly, the matrix $B(\mathbf{k})$ takes the following form

$$B(\mathbf{k}) = \begin{bmatrix} B_1(\mathbf{k}) & & & & \\ & B_2(\mathbf{k}) & & & \\ & & B_3(\mathbf{k}) & & \\ & & & \ddots & \\ & & & & B_N(\mathbf{k}) \end{bmatrix} \quad (16)$$

where each $B_m(\mathbf{k})$ is an $U(2)$ matrix. Therefore, the map f splits into N maps

$$f_m : T^3 \rightarrow U(2) \quad m = 1, 2, \dots, N \quad (17)$$

The next step is reducing $U(2)$ to $SU(2)$, which is a three-dimensional manifold. The motivation of this step will be clear later when we consider the degree of certain map. To this end, we factorize $B_m(\mathbf{k}) = \exp[i\theta_m(\mathbf{k})]u_m(\mathbf{k})$, where $u_m(\mathbf{k})$ is a $SU(2)$ matrix function of $\mathbf{k}$. Because of the facts that $\exp[i\theta_m(\mathbf{k})]u_m(\mathbf{k}) = -\exp[i\theta_m(\mathbf{k})][-u_m(\mathbf{k})]$ and $-u_m(\mathbf{k})$ is also $SU(2)$ matrix, there is ambiguity in this factorization. To avoid this ambiguity, we can choose a point $\mathbf{k}_0$ in T^3 , and choose one factorization $B_m(\mathbf{k}_0) = \exp[i\theta_m(\mathbf{k}_0)]u_m(\mathbf{k}_0)$. The $U(1)$ factors of other $\mathbf{k}$ points are determined by

$$\theta_m(\mathbf{k}) = \theta_m(\mathbf{k}_0) - \frac{i}{2} \int_{\mathbf{k}_0}^{\mathbf{k}} d\mathbf{k} \nabla_{\mathbf{k}} \det[B_m(\mathbf{k})] \quad (18)$$

It is easy to check that with this equation $u_m(k)$ will have unitary determinant. To make the integral in the above equation unambiguous, we require that $\oint_l d\mathbf{k} \nabla_{\mathbf{k}} \det[B_m(\mathbf{k})] = 0$ for arbitrary loop l in T^3 . For contractable loops, this is trivially satisfied. For non-contractable loop, consider the non-contractable loop AOB in Fig.2 as an example, we have




Figure 2. The loop (closed line) AOB in T^3 . The point O is $k = (0, 0, 0)$. Points A and B are the same point because the periodic boundary condition. AOB is homotopically nontrivial, *i.e.* AOB cannot be continuously deformed to a single point.

$$\oint_{AOB} d\mathbf{k} \nabla_{\mathbf{k}} \det[B_m(\mathbf{k})] = \int_{AO} d\mathbf{k} \nabla_{\mathbf{k}} \det[B_m(\mathbf{k})] + \int_{OB} d\mathbf{k} \nabla_{\mathbf{k}} \det[B_m(\mathbf{k})] = 0 \quad (19)$$

which is a result of the relation $\det[B_m(-\mathbf{k})] = \det[B_m(\mathbf{k})]$. Therefore, we have a factorization of $U(2)$ into $SU(2)$ and $U(1)$. With this factorization, we also have that $\oint_l d\mathbf{k} \nabla_{\mathbf{k}} \theta_m(\mathbf{k}) = 0$ (for arbitrary loop l). Therefore, we can adiabatically deform the $U(1)$ factor to 1. Having got rid of the $U(1)$ factor, the maps $f_m : T^3 \rightarrow U(2)$ are deformed to $g_m : T^3 \rightarrow SU(2)$. This completes our discussion on matrix $B(\mathbf{k})$. With the aid of $B(\mathbf{k})$, we can give geometrical interpretation of topological invariants of TR invariant topological insulators.

In order to prove the equivalence, we first show how to relate the topological invariant P_3 in Eq. (4) to a winding number. Similar formula appeared in Ref.[12] in a slightly different form but we would like to include the discussion here to make our discussion self-contained. We start from the integral topological invariants

$$P_3 = \frac{1}{16\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr}\{[f_{ij}(\mathbf{k}) - \frac{2}{3}ia_i(\mathbf{k}) \cdot a_j(\mathbf{k})] \cdot a_k(\mathbf{k})\} \quad (20)$$

which is valid independent of the TR symmetry. In a TR invariant system, with the aid of $B(\mathbf{k})$, we have

$$\begin{aligned}
a_i^{\alpha\beta}(-\mathbf{k}) &= -i\langle -\mathbf{k}, \alpha | \partial_{-\mathbf{k}_i} | -\mathbf{k}, \beta \rangle \\
&= i \sum_{\alpha'\beta'} B_{\alpha\alpha'} \langle \Theta, \mathbf{k}, \alpha' | \partial_{\mathbf{k}_i} (B_{\beta\beta'}^* | \Theta, \mathbf{k}, \beta' \rangle) \\
&= i \sum_{\alpha'\beta'} B_{\alpha\alpha'} B_{\beta\beta'}^* \langle \Theta, \mathbf{k}, \alpha' | \partial_{\mathbf{k}_i} | \Theta, \mathbf{k}, \beta' \rangle + i \sum_{\alpha'\beta'} B_{\alpha\alpha'} \partial_{\mathbf{k}_i} B_{\beta\beta'}^* \delta_{\alpha'\beta'} \\
&= \sum_{\alpha'\beta'} B_{\alpha\alpha'} (a_i^{\alpha'\beta'}(\mathbf{k}))^* B_{\beta'\beta}^\dagger + i \sum_{\alpha'} B_{\alpha\alpha'} \partial_{\mathbf{k}_i} B_{\alpha'\beta}^\dagger
\end{aligned} \tag{21}$$

which can be simply written as

$$a_i(-\mathbf{k}) = B(\mathbf{k}) a_i^*(\mathbf{k}) B^\dagger(\mathbf{k}) + i B(\mathbf{k}) \partial_i B^\dagger(\mathbf{k}) \tag{22}$$

Therefore, the field strength satisfies the following relation

$$\begin{aligned}
f_{ij}(-\mathbf{k}) &= \partial_{-\mathbf{k}_i} a_j(-\mathbf{k}) - \partial_{-\mathbf{k}_j} a_i(-\mathbf{k}) + i[a_i(-\mathbf{k}), a_j(-\mathbf{k})] \\
&= -B(\mathbf{k}) f_{ij}^*(\mathbf{k}) B^\dagger(\mathbf{k})
\end{aligned} \tag{23}$$

These results imply that the non-abelian Berry potential and curvature at $-\mathbf{k}$ are simply related to those at $\mathbf{k}$ by a non-abelian gauge transformation $B(\mathbf{k})$. Therefore, we obtain that

$$\begin{aligned}
P_3 &= \frac{1}{16\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} \{ [f_{ij}(-\mathbf{k}) - \frac{2}{3} i a_i(-\mathbf{k}) a_j(-\mathbf{k})] a_k(-\mathbf{k}) \} \\
&= -\frac{1}{16\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} \{ [f_{ij}(\mathbf{k}) - \frac{2}{3} i a_i(\mathbf{k}) a_j(\mathbf{k})] a_k(\mathbf{k}) \}^* \\
&\quad - \frac{i}{8\pi} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} \partial_i (B a_j^* \partial_k B^\dagger) \\
&\quad - \frac{1}{24\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} [(B \partial_i B^\dagger)(B \partial_j B^\dagger)(B \partial_k B^\dagger)] \\
&= -P_3^* - \frac{1}{24\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} [(B \partial_i B^\dagger)(B \partial_j B^\dagger)(B \partial_k B^\dagger)] \\
&= -P_3 - \frac{1}{24\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} [(B \partial_i B^\dagger)(B \partial_j B^\dagger)(B \partial_k B^\dagger)]
\end{aligned} \tag{24}$$

or

$$2P_3 = -\frac{1}{24\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} [(B \partial_i B^\dagger)(B \partial_j B^\dagger)(B \partial_k B^\dagger)] \tag{25}$$

Both the LHS and RHS of the above equation depend on gauge choices, but the parity of $2P_3$ is gauge invariant

$$2P_3(\text{mod } 2) = -\frac{1}{24\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr} [(B \partial_i B^\dagger)(B \partial_j B^\dagger)(B \partial_k B^\dagger)] (\text{mod } 2) \tag{26}$$

which is the important quantity for characterization of the topological insulator.

In our previous discussion, we have split $B(k)$ into direct sum of $SU(2)$ matrices $(B_m, m = 1, 2, \dots, N)$. It is readily obtained that

$$2P_3(\text{mod } 2) = \sum_{m=1}^N \nu_m \pmod{2} \tag{27}$$

where

$$\nu_m = -\frac{1}{24\pi^2} \int d^3\mathbf{k} \epsilon^{ijk} \text{Tr}[(B_m \partial_i B_m^\dagger)(B_m \partial_j B_m^\dagger)(B_m \partial_k B_m^\dagger)] \quad (28)$$

We notice that ν_m is just the winding number, or the integral form of degree of the map

$$g_m : T^3 \rightarrow SU(2) \quad (29)$$

The mod 2 degree of this map is given by

$$\deg_2(g_m) = \nu_m \pmod{2} \quad (30)$$

Therefore we have

$$2P_3 \pmod{2} = \sum_{m=1}^N \deg_2(g_m) \quad (31)$$

This is the integral form of degree of map. The natural question is whether we can find a discrete form, as discussed in the previous section. Let us denote the image of $\mathbf{k}$ under the map g_m as $B_m(\mathbf{k})$, which is a $SU(2)$ matrix. As has been noted, $B_m(\Gamma_i)$ is anti-symmetric at TRIM Γ_i . There are only two anti-symmetric matrices in $SU(2)$, given by

$$A_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

It is easy to see that $\text{Pf}[A_1] = 1$ and $\text{Pf}[A_2] = -1$. In principle, we can pick any image point on the $SU(2)$ to perform the counting for the degree of the map. However, because of the TR symmetry, the counting is particularly simple if we pick either A_2 or A_1 . We now count the number of source points $\mathbf{k}$, modulo 2, which map onto A_2 . We first notice that if $B_m(\mathbf{k}) = A_2$, then $B_m(-\mathbf{k}) = -B_m^T(\mathbf{k}) = A_2$. Therefore, if $\mathbf{k}$ maps to A_2 , so does $-\mathbf{k}$. If $\mathbf{k}$ is not one of the TRIM, $\mathbf{k}$ and $-\mathbf{k}$ are distinct, and these two points do not contribute to $\deg_2(g_m)$, which counts the number of source points modulo 2. The only source points which may contribute to $\deg_2(g_m)$ are TRIM, where $\mathbf{k}$ and $-\mathbf{k}$ are equivalent. It is also important to note that the calculation of mod 2 degree of map is easier than that of the integer degree, which involves the $\pm$ signs of the Jacobian of the map. Because $-1 = 1 \pmod{2}$, we do not need to be concerned about the signs of the Jacobian when calculating degree of map mod 2.

Because we have reduced the map from $U(2)$ to $SU(2)$, $\det[B(\Gamma_i)] = 1$, we have $\delta_i = \sqrt{\det[B(\Gamma_i)]} / \text{Pf}[B(\Gamma_i)] = \text{Pf}[B(\Gamma_i)] = \prod_m \text{Pf}[B_m(\Gamma_i)]$. Suppose that there are n_m TRIMs which map onto A_2 under the map g_j and $8 - n_m$ TRIMs which map onto A_1 . By counting the number of points which map to A_2 mod 2, we have $\deg_2(g_m) = n_m \pmod{2}$, and therefore

$$(-1)^{2P_3} = (-1)^{\sum_{m=1}^N \deg_2(g_m)} = \prod_m (-1)^{n_m} \quad (32)$$

On the other hand, the discrete invariant defined in Ref. [10] is given by

$$(-1)^{\nu_0} = \prod_i \text{Pf}[B(\Gamma_i)] = \prod_{i,m} \text{Pf}[B_m(\Gamma_i)] = \prod_m (-1)^{n_m} \quad (33)$$

Therefore, we proved the central result of this paper, namely the exact equivalence between the integral invariant of QHZ and the discrete invariant of FKM for the 3D topological insulator:

$$(-1)^{2P_3} = (-1)^{\nu_0} \quad (34)$$

In the proof given above, we made the assumption that no accidental degeneracy except the Kramers degeneracy occurs, which simplified the discussion. The extension to the generic case is straightforward because $\pi_3(U(2N)) = \pi_3(U(2))$ holds for any integer $N \geq 1$. One can always deform the map f to one of the $U(2)$ subgroups of $U(2N)$, so that the proof discussed above applies.

Acknowledgments

We thank Shao-Long Wan and Yong-Shi Wu for helpful discussions. This work is supported by the US Department of Energy, Office of Basic Energy Sciences under contract DE-AC03-76SF00515. Z. Wang acknowledges the support of China Scholarship Council, NSF of China(Grant No.10675108).

Appendix A. Discussion on the global basis of wavefunctions

In this appendix we shall show that the global basis of wavefunction exists on T^3 at presence of time-reversal symmetry. Because the global definition of $B(k)$ and the discrete invariant ν_0 depends on the existence of such global basis, this point is important to our argument.

We divide the $2N$ filled bands into N TR pairs. Within each pair, the Hilbert space is two dimensional at each momentum k . Therefore, the Hilbert space is naturally an $U(2)$ fibre bundle on T^3 . Generically it is not evident that this bundle is trivial. The key to the existence of global basis is the TR symmetry.

The “surface” of the cubic in Fig.2 consists of three T^2 (two-dimensional tori). We consider the restriction of the $U(2)$ bundle to one of these tori. The bundle on T^2 is trivial because the first chern number $C_1 = 0$, which is a consequence of TR symmetry. Therefore, a global basis exists on the surface of the cubic. Such a basis can be extrapolated to the interior of the cubic since the interior region is topologically trivial. We consider the transition function between the surface and the interior region on their overlapping region, which is topologically equivalent to S^2 (two-dimensional sphere). Because the second homotopy group $\pi_2(U(2))$ is trivial, mapping from S^2 to $U(2)$ (the structure group of the fibre bundle) are all trivial. Thus the bundle is trivial on T^3 . Therefore, global basis does exist.

For comparison, we note that in the case of integer quantum Hall states, the global basis does not exist on the Brillouin zone T^2 , because the relevant fibre bundle on T^2 is non-trivial. From this example we also see that TR symmetry is necessary for our argument.

References

- [1] Zhang S C 1992 *Int. J. Mod. Phys. B* **6** 25
- [2] Thouless D J, Kohmoto M, Nightingale M P and den Nijs M 1982 *Phys. Rev. Lett.* **49** 405
- [3] B A Bernevig, T L Hughes and S C Zhang 2006 *Science* **314** 1757
- [4] König M, Wiedmann S, Brüne C, Roth A, Buhmann H, Molenkamp L, Qi X L and Zhang S C 2007 *Science* **318** 766–770
- [5] Fu L and Kane C L 2007 *Phys. Rev. B* **76** 045302
- [6] Hsieh D, Qian D, Wray L, Xia Y, Hor Y S, Cava R J and Hasan M Z 2008 *Nature* **452** 970–974
- [7] Zhang H, Liu C X, Qi X L, Dai X, Fang Z and Zhang S C 2009 *Nature Physics* **5** 438 – 442
- [8] Xia Y, Qian D, Hsieh D, Wray L, Pal A, Lin H, Bansil A, Grauer D, Hor Y S, Cava R J and Hasan M Z 2009 *Nat. Phys.* **5** 398
- [9] Chen Y L, Analytis J G, Chu J H, Liu Z K, Mo S K, Qi X L, Zhang H J, Lu D H, Dai X, Fang Z, Zhang S C, Fisher I R, Hussain Z and Shen Z X 2009 *Science* **325**
- [10] Fu L, Kane C L and Mele E J 2007 *Phys. Rev. Lett.* **98** 106803
- [11] Murakami S, Nagaosa N and S C Zhang 2003 **301** 1348
- [12] Qi X L, Hughes T and Zhang S C 2008 *Phys. Rev. B* **78** 195424
- [13] Moore J E and Balents L 2007 *Phys. Rev. B* **75** 121306
- [14] F. Wilczek, *Nature* **458**, 129 (2009)
- [15] Qi X L, Li R, Zang J and Zhang S C 2009 *Science* **323** 1184–1187
- [16] Zhang S C and Hu J P 2001 *Science* **294** 823
- [17] Bernevig B A, Chern C H, Hu J P, Toumbas N and Zhang S C 2002 *Annals of Physics* **300** 185
- [18] Fu L and Kane C L 2006 *Phys. Rev. B* **74** 195312
- [19] Essin A M, Moore J E and Vanderbilt D 2009 *Phys. Rev. Lett.* **102** 146805
- [20] Dubrovin B A, Fomenko A T and Novikov S P 1985 *Modern Geometry—Methods and Applications Part 2: The Geometry and Topology of Manifolds* (Springer)